



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE, BY E. B. SEITZ.—The value of g , in Mr. Baker's solution, p. 25, may be found as follows:

Let $g = u_n$; then $u_n = u_{n-1} \pm 1 \dots (1)$, and $u_{n-1} = u_{n-2} \mp 1, \dots (2)$ the upper sign being used when n is odd, and the lower when n is even. Adding (1) and (2), we have $u_n - u_{n-1} - 2u_{n-2} = 0$, an equation in Finite Differences, whose solution gives $u_n = C_1 \cdot 2^n + C_2(-1)^n \dots (3)$

When $n = 1$, $u_1 = 2C_1 - C_2 = 1 \dots (4)$, and when $n = 2$, $u_n = 4C_1 + C_2 = 1 \dots (5)$. From (4) and (5) we find $C_1 = \frac{1}{3}$, and $C_2 = -\frac{1}{3}$; therefore $g = u_n = \frac{1}{3}(2^n \pm 1)$, the double sign being used as above. Hence the angles of the n th triangle are

$$\frac{1}{3}\pi \pm (\frac{1}{2})^n(\frac{1}{3}\pi - A), \quad \frac{1}{3}\pi \pm (\frac{1}{2})^n(\frac{1}{3}\pi - B), \quad \frac{1}{3}\pi \pm (\frac{1}{2})^n(\frac{1}{3}\pi - C).$$

[Mr. Baker has sent a "Revised Solution" of the question, in which he represents g by the same formula obtained above, and also notices the error in the table, pointed out by Prof. Johnson, but our space will not permit its publication.]

ANSWER TO QUERY (SEE P. 176, VOL. IV), BY THE EDITOR.

FROM the fixed point E draw a line EAB perpendicular to the fixed line CI and intersecting it in B , at any distance a from the point E , and let A represent the middle point of EB .

Let FGH be any position of the right angle, the side GH intersecting the line CI in L , and let the fix'd length of $GE = a$; then, to find the locus of P , the middle point of GF , draw PQ perpendicular to AB , intersecting the side GH in m ; draw Gn perpendicular to CI , and put $AQ = x$ and $PQ = y$.

Because $AE = \frac{1}{2}a$, $EQ = \frac{1}{2}a + x$, and $BQ = \frac{1}{2}a - a + x = x - \frac{1}{2}a$; $\therefore Gn = 2(x - \frac{1}{2}a)$ and $Fn = \sqrt{a^2 - [2(x - \frac{1}{2}a)]^2} = \pm 2\sqrt{(x^2 - ax)}$.

Hence, from the similar triangles FnG and EQM , and FnG and PGm ,

$$Fn : Gn :: EQ : Qm, \text{ or } 2\sqrt{(x^2 - ax)} : 2(x - \frac{1}{2}a) :: x + \frac{1}{2}a : Qm = \frac{x^2 - \frac{1}{4}a^2}{\sqrt{(x^2 - ax)}},$$

$$Fn : FG :: PG : Pm, \text{ or } 2\sqrt{(x^2 - ax)} : a :: \frac{1}{2}a : Pm = \frac{\frac{1}{4}a^2}{\sqrt{(x^2 - ax)}}.$$

$$\text{But } Qm + Pm = y = \frac{x^2 - \frac{1}{4}a^2}{\sqrt{(x^2 - ax)}} + \frac{a^2}{\sqrt{(x^2 - ax)}} = \frac{x^2}{\sqrt{(x^2 - ax)}}; \therefore y^2 = \frac{x^3}{x - a},$$

which is the equation to the cissoid.



